

Long-Term Variable Milfoil Management and Control Plan for GLEN LAKE Goffstown, New Hampshire Hillsborough County

Prepared by: New Hampshire Department of Environmental Services (DES),
in consultation with the
New Hampshire Fish and Game Department (F&G)
Prepared: February 2008

PROBLEM STATEMENT

Exotic aquatic plants pose a threat to the ecological, aesthetic, recreational, and economic values of lakes and ponds (Luken & Thieret, 1997, Halstead, 2000). According to the 2006 Section 305(b) and 303(d) Consolidated Assessment and Listing Methodology (CALM), “exotic macrophytes are non-native, fast growing aquatic plants, which can quickly dominate and choke out native aquatic plant growth in the surface water. Such infestations are in violation of Env-Ws 1703.19, which states that surface waters shall support and maintain a balanced, integrated and adaptive community of organisms having a species composition, diversity, and functional organization comparable to that of similar natural habitats of a region” (DES, 2006).

Though exotic aquatic plants can negatively impact an aquatic system, native aquatic plants are beneficial to the aquatic ecology of waterbodies. Diverse assemblages of native aquatic plants are a source of oxygen to the system, they provide stabilizing root systems to minimize erosion and turbidity, and they provide food and habitat for aquatic life.

Variable milfoil (*Myriophyllum heterophyllum*) was first documented in Glen Lake in Goffstown, New Hampshire in 2007, though we expect the new infestation of this waterbody was in 2006.

Figure 1 illustrates the distribution of variable milfoil infestations in this waterbody, as of fall 2007. At this time, only one small area (2.38 acres) of variable milfoil was found during the plant mapping activities. The milfoil in this area ranges from roughly 65% to 90% cover.

In terms of the impacts of the variable milfoil in the system, there are several houses around the shoreline of Glen Lake, with mostly year round homes. There are also numerous back lots with lake rights via designated rights of ways going down to the lake edge.

Lake residents have expressed concerns about variable milfoil spreading further throughout the lake system to a level that could ultimately impede recreational uses of the waterbody. Additional concerns are that variable milfoil fragments could spread down through the Piscataquog River system and increase areas of milfoil infestation in the Merrimack River downstream.

At this time, there are no data and no observed problems with the biological integrity of the aquatic community as a result of the variable milfoil infestation; however, the variable milfoil

infestation is still somewhat localized. No biological integrity surveys have been conducted, however, as part of this plan preparation.

PURPOSE

The purposes of this exotic aquatic plant management and control plan are:

1. To identify the waterbody's beneficial use areas, including essential aquatic habitat, designated conservation zones, swimming areas, boat access sites, and boating use areas;
2. To present the aquatic macrophyte distribution map, including both native and exotic species;
3. To identify short-term and long-term exotic aquatic plant control goals that protect and conserve the lake's beneficial uses;
4. To recommend exotic plant control actions that meet the goals outlined in this plan; and
5. To recommend monitoring strategies to determine the success of the control practices over time in meeting the goals.

This plan also summarizes the current physical, biological, ecological, and chemical components of Glen Lake and the social and ecological impacts of the milfoil infestation. The intent of this strategic plan is to eradicate variable milfoil from Glen Lake over time through the use of Integrated Pest Management Strategies (IPM). Appendix A details the strategies available for waterbodies with exotic species, and provides more information on each of the activities that are recommended within this plan.

GOALS/OBJECTIVES OF MILFOIL CONTROL ACTIONS

The goal for Glen Lake is the eventual eradication of variable milfoil from the system using an Integrated Pest Management Approach. To achieve this goal, we recommend the following:

- 1) In spring 2008, conduct a 2,4-D treatment to reduce the size and density of the variable milfoil patch in Glen Lake.
- 2) In summer 2008 and annually thereafter, monitor the lake for variable milfoil re-growth or new growth and manage it using non-chemical approaches.

DES also recommends that the lake residents maintain a Weed Watcher program and Lake Host Program or other educational effort at the public access of the pond.

Town Support

As this is a new infestation, DES did not seek additional funding from the town or the lake residents. The town has been very helpful in providing information about abutters to Glen Lake.

Glen Lake Residents

At this time there is no organized lake association on Glen Lake. There is a group called the Goffstown Residents Association that has a website where local information and lake information is posted regularly for the community to view, and it is updated regularly by one lake resident.

In discussions with representatives of the lake, they are supportive of the efforts to control variable milfoil, and are very interested in starting up a Weed Watcher Program in the spring/summer of 2008 to help to monitor the waterbody for new milfoil growths.

The lake is actively involved in the DES Volunteer Lake Assessment Program (VLAP), and there is an obvious connection between lake residents to protect Glen Lake.

WATERBODY CHARACTERISTICS

The following table summarizes basic physical and biological characteristics of Glen Lake.

General Lake Information	
Lake area (acres)	118.8
Watershed area (acres)	129,428.0
Shoreline Uses (residential, forested, agriculture)	Residential, forested
Max Depth (ft)	52.1
Mean Depth (ft)	19.5
Trophic Status	Mesotrophic
Color (CPU) in Epilimnion	22
Clarity (ft)	5.9
Flushing Rate (yr ⁻¹)	80.0
Natural waterbody/Raised by Damming/Other	Artificial
Plant Community Information Relative to Management	
Invasive Plants (Latin name)	<i>Myriophyllum heterophyllum</i> , <i>European naiad</i> , <i>purple loosestrife</i>
Infested Area (acres)	Approximately 2.38 acres with variable milfoil
Distribution (ringing lake, patchy growth, etc)	Dense localized area of infestation denoted in Figure 1.
Sediment type in infested area (sand/silt/organic/rock)	Sandy/rocky/silty
Rare, Threatened, or Endangered Species in Waterbody (according to NH Natural Heritage Inventory)	Brook Floater- State Endangered Wood Turtle-Rare in NH
Area of Littoral Zone (acres)	53
Area of Profundal Zone (acres)	84
Area of Macrophyte Coverage (native or otherwise) of Plants in Littoral Zone	26
% of Littoral Zone with Macrophyte Cover	50
% of Macrophyte cover comprised of invasives	9
% of Littoral Zone with Variable Milfoil Cover	5

An aquatic vegetation map and key from an August 21, 2007 survey by the DES Biology Section is shown in Figure 2. A bathymetric map is shown in Figure 3.

BENEFICIAL (DESIGNATED) USES

In New Hampshire, beneficial (designated) uses of our waterbodies are grouped into five general categories: Aquatic Life, Fish Consumption, Recreation, Drinking Water Supply, and Wildlife (CALM).

Of these, Aquatic Life and Recreation are the ones affected by the presence of invasive plants like variable milfoil.

AQUATIC LIFE

The goal for aquatic life support is to provide suitable chemical and physical conditions for supporting a balanced, integrated and adaptive community of aquatic organisms having a species composition, diversity, and functional organization comparable to that of similar natural habitats of the region.

FISHERIES AND WILDLIFE

Glen Lake is a man-made lake that was created by damming the Piscataquog River. There is excellent bottom structure in the form of submerged trees and stumps on one side, while the opposite side exhibits a rocky shoreline with a rapid drop-off. Fish species present include largemouth bass, smallmouth bass, golden shiner, pickerel, yellow perch, white perch, common white sucker, brown bullhead, and yellow bullhead. Brook trout and rainbow trout are also stocked each year. Fishing pressure is light during open-water and rare during winter.

Figure 4 shows the common fishing locations on Glen Lake.

A Natural Heritage Inventory review showed that the brook floater mussel and the wood turtle have been recorded in or near Glen Lake. The brook floater mussel is listed as state endangered, and the wood turtle is listed as rare in New Hampshire. Because this treatment and additional non-chemical control practices affect only a very small portion of this waterbody (2.38 acres out of a total 119 acres, impacts to either of these species is unlikely, as ample habitat and refugia will be present during and after the management practices.

RECREATIONAL USES AND ACCESS POINTS

Glen Lake is used for numerous recreational activities, including boating, fishing, swimming, and water skiing by both lakefront residents and transient boaters. There is one designated public access for boats on the northeastern side of the pond. Motor boats, as well as kayaks and canoes can use this facility. There is ample parking for vehicles with trailers across the street from the ramp. Figure 5 illustrates the typical boat paths for the pond.

There is one picnic area with a sandy beach near the public access site. There are a few small private swim beaches located on private properties around the pond. Figure 6 shows the locations commonly used for swimming, and the locations of swim platforms and docks on Glen Lake.

MACROPHYTE EVALUATION

The littoral zone is defined as the nearshore areas of a waterbody where sunlight penetrates to the bottom sediments. The littoral zone is typically the zone of rooted macrophyte growth in a waterbody.

The littoral zone of Glen Lake is characterized by a mix of native and non-native (variable milfoil and European naiad) plant growth (Figure 2). Native species include a mix of floating plants (floating heart, watershield, duckweed), emergent plants (grassy arrowhead, pipewort), and submergent plants (water naiad, tapegrass, muskgrass, pondweeds). Native plant communities are mixed around the entire lake, and are characterized as ‘scattered’ by the DES.

There are no records of state threatened or endangered plant species.

The invasive plants, European naiad and purple loosestrife, were also found in and around Glen Lake. European naiad was present in one cover on the south-central side of Glen Lake. It was scattered in nature, and can be removed by simple hand-removal and bottom barrier placement, which DES will coordinate in summer 2008. DES will work with the lake residents to educate them about the purple loosestrife, and how to effectively hand-remove it.

HISTORICAL CONTROL ACTIVITIES ON THIS WATERBODY:

This is a new infestation. No previous control measures have been documented.

MILFOIL MANAGEMENT OPTIONS

The control practices used should be as specific to milfoil as feasible. No control of native aquatic plants is intended.

Exotic aquatic plant management relies on a combination of proven methods that control exotic plant infestations, including physical control, chemical control, biological controls (where they exist), and habitat manipulation. Integrated Pest Management Strategies (IPM) are typically implemented using Best Management Practices (BMPs) based on site-specific conditions so as to maximize the long-term effectiveness of control strategies. Descriptions for the control activities are closely modeled after those prescribed by the Aquatic Ecosystem Restoration Foundation (AERF) (2004). This publication can be found online at http://www.aquatics.org/aquatic_bmp.pdf.

Criteria for the selection of control techniques are presented in Appendix A. Appendix B includes a summary of the exotic aquatic plant control practices used by the State of New Hampshire. DES has evaluated the feasibility of potential control practices on Glen Lake. The following table summarizes DES’ control strategy recommendations for Glen Lake.

FEASIBILITY EVALUATION FOR CONTROL ALTERNATIVES

Control Method	Use on Glen Lake
Restricted Use Areas	A Restricted Use Area, or at the very least a fragment barrier, are recommended for use in containing this isolated patch of variable milfoil. DES will establish a containment area on Glen Lake in summer 2008, following the herbicide treatment, as needed.

Control Method	Use on Glen Lake
Hand-pulling	DES recommends hand-removal of persistent fragments or new growth of variable milfoil following the herbicide treatment. DES also recommends that the European naiad and the purple loosestrife be controlled with hand-removal as well. In an effort to both educate lake residents about the exotic plants, and to garner more local support in early detection efforts, DES will work with representatives from the lake to establish a Weed Watcher Program to facilitate more hand-removal (non-chemical) control options in the future.
Mechanical Harvesting/Removal	For Glen Lake, mechanical harvesting is not recommended due to the threat of spreading variable milfoil to uninfested areas of the lake through the generation of fragments.
Benthic Barriers	For Glen Lake, DES recommends installing small benthic barriers in areas of re-growth if small patches of variable milfoil re-grow and can adequately be contained by benthic barriers. Benthic barriers may also be used for the European naiad.
Herbicides	For Glen Lake, herbicide use is recommended as primary treatment due to percent cover of the variable milfoil over the 2.38 acres of infested lake bottom. The goal is to reduce the density of variable milfoil and its rooting systems, to better facilitate hand-removal and other non-chemical approaches.
Extended Drawdown	Drawdown is not an effective control method for variable milfoil.
Dredge	Not recommended due to nature of exotic plant distribution, the cost, or the ancillary ecological impacts that the dredge could have.
Biological Control	There are no approved biological controls for variable milfoil at this time in New Hampshire.
No Control	This is a small new infestation of variable milfoil, and as such, it warrants a rapid response at management. If left unmanaged, the variable milfoil has the great potential to spread along most of the northern (and shallower) shoreline of Glen Lake.

EXOTIC AQUATIC PLANT CONTROL PLAN

An evaluation of the size, location, and type of variable milfoil infestation, as well as the waterbody uses was conducted by DES during September 21, 2007. Based on the evaluation, the following control actions are recommended:

Year	Treatment Type	Responsible Party	Schedule
2008	2,4-D treatment (or Diquat as an alternative) of variable milfoil in the area shown in Figure 1	Aquatic Control Technology, Inc.	May/June
	Train local residents to act as Weed Watchers around Glen Lake	DES	Late June or early July

Year	Treatment Type	Responsible Party	Schedule
	Hand-removal of variable milfoil stems after herbicide treatment, and removal of European naiad. Benthic barriers may also be used.	DES divers or contract divers	July through September
	Purple loosestrife non-herbicide control	Local residents	Summer months
	Educational material posted at launch/access	Local residents	June
	Site assessment	DES	August/September
2009	Hand-removal, diver assisted suction harvesting, or benthic barrier placement to control variable milfoil and European naiad	DES	Summer/Fall
	Weed Watcher monitoring	Local residents	June through September
	Access site education information posting	Local residents	June through September
	Site assessment	DES	August/September
2010	2,4-D treatment (or Diquat as an alternative) of variable milfoil only if needed (based on fall 2009 assessment)	TBD	May/June
	Hand-removal, diver assisted suction harvesting, or benthic barrier placement to control variable milfoil and European naiad	DES	Summer/Fall
	Weed Watcher monitoring	Local residents	June through September
	Access site education information posting	Local residents	June through September
	Site assessment	DES	August/September
2011	Hand-removal, diver assisted suction harvesting, or benthic barrier placement to control variable milfoil and European naiad	DES	Summer/Fall
	Weed Watcher monitoring	Local residents	June through September
	Access site education information posting	Local residents	June through September
	Site assessment	DES	August/September

Year	Treatment Type	Responsible Party	Schedule
2012	Hand-removal, diver assisted suction harvesting, or benthic barrier placement to control variable milfoil and European naiad	DES	Summer/Fall
	Weed Watcher monitoring	Local residents	June through September
	Access site education information posting	Local residents	June through September
	Site assessment	DES	August/September
2013	Management Plan Update	DES and interested parties	Fall

- Approximately 2.38 acres of the waterbody will be targeted for the herbicide treatment (approximately 2% of the surface area).
- The Department of Agriculture will impose standard short-term use restrictions for specified days depending on the use (irrigation, contact, etc) and the herbicide used. The shoreline will be posted and public notice will be made.
- By recommending follow-up management practices that utilize integrated plant management strategies such as benthic barrier placement and hand-pulling re-growth, variable milfoil re-growth or population expansion can be slowed.
- Based on the types of native plants that are mixed in with the stands of variable milfoil (Figure 2) where herbicide application is recommended there are no significant impacts to native plant communities. It is expected that a well distributed stand of native aquatic plants will remain following herbicide application.
- It is important to realize that aquatic herbicide applications are conducted in a specific and scientific manner, and that the herbicides that are used can be target-specific when used at appropriate doses/concentrations: this means that the invasive plant can be removed and native plants favored in this type of control practice. *Not all aquatic plants will be impacted as a result of an herbicide treatment.*
- Because this is a natural system that is being evaluated for management, it is impossible to accurately predict a management course over five years that could be heavily dependent on uncontrolled natural circumstances (weather patterns, temperature, etc). This management plan should be considered a dynamic document that is geared to the actual field conditions that present themselves in this waterbody. If circumstances arise that require the modification of part or all of the recommendations outline here, all interested parties will be consulted for their input on revisions that may be needed to further the goal of variable milfoil management in the subject waterbody.

Figure 1- Map of Milfoil Infestation





Figure 2- Aquatic Vegetation Map and Key



Symbol	Common Name	Latin Name
N	European naiad	<i>Najas minor</i>
J	Native naiad	<i>Najas sp.</i>
V	Tapegrass	<i>Vallisneria americana</i>
G	Grassy arrowhead	<i>Sagittaria graminea</i>
L	Duckweed	<i>Lemna</i>
3	Muskgrass	<i>Chara</i>
4	Purple loosestrife	<i>Lythrum salicaria</i>
B	Watershield	<i>Brasenia schreberi</i>
E	Pipewort	<i>Eriocaulon</i>
X	Pondweed	<i>Potamogeton sp.</i>
H	Floating heart	<i>Nymphoides cordata</i>
M	Variable milfoil	<i>Myriophyllum heterophyllum</i>

Figure 3- Bathymetric Map of Glen Lake, Goffstown




Figure 4- Common Fishing Locations




Figure 5- Common Boating Lanes





Figure 6- Swim Areas, Docks, and Swim Rafts



APPENDIX A

CRITERIA TO EVALUATE THE SELECTION OF AQUATIC PLANT CONTROL TECHNIQUES

Preliminary Investigations

I. Field Site Inspection

- Verify genus and species of the plant.
- Determine if the plant is a native or exotic species per RSA 487:16, II.
- Map extent of the plant infestation (area, water depth, height of the plant, density of the population).
- Document any native plant abundances and community structure around and dispersed within the exotic/nuisance plant population.

II. Office/Laboratory Research of Waterbody Characteristics

- Contact the appropriate agencies to determine the presence of rare or endangered species in the waterbody or its prime wetlands.
- Determine the basic relevant limnological characteristics of the waterbody (size, bathymetry, flushing rate, nutrient levels, trophic status, and type and extent of adjacent wetlands).
- Determine the potential impacts to downstream waterbodies based on limnological characteristics (water chemistry, quantity, quality).

Overall Control Options

For any given waterbody that has an infestation of exotic plants, one of three options will be selected, based on the status of the infestation, the available management options, and the technical knowledge of the DES Limnologists who have conducted the field work and who are preparing this plan. The options are as follows:

- 1) **Eradication:** Herbicide application targeted at exotic aquatic plant to be eradicated, to either eradicate the plant or to reduce overall biomass to a point where alternative non-chemical strategies may be used. This action will be followed by thorough annual monitoring for regrowth and the use of non-chemical actions to achieve the eradication.
- 2) **Containment:** The aim of this approach is to limit the size and extent of the existing infestation. An herbicide application may be used to reduce specified areas down to a percent cover of the exotic species so that it can be maintain or contained with alternative management strategies, including Restricted Use Areas, benthic barriers, and others. Subsequent herbicide applications may be necessary if the target species shows exponential growth and further spread.

- 3) No action. If the infestation is too large, spreading too quickly, and past management strategies have proven ineffective at controlling the target exotic aquatic plant, DES, in consultation with others, may elect to recommend ‘no action’ at a particular site. All efforts will instead be made towards containment of the target species to that specific waterbody, so that downstream migration of the plant can be prevented.

If eradication or control is the recommended option to pursue, the following series of control techniques may be employed. The most appropriate technique based on the determinations of the preliminary investigation will be selected.

Guidelines and requirements of each control practice are detailed below each alternative.

A. Hand-Pulling

- Can be used for exotic or native species.
- Can be used if infestation is in a small localized area (sparsely populated patch of up to 5' X 5', single stems, or dense small patch up to 2' X 2').
- Can be used if plant density is low, or if target plant is scattered and not dense.
- Can be used if the plant could effectively be managed or eradicated by hand-pulling a few scattered plants.
- Use must be in compliance with the Wetlands Bureau rules.

B. Mechanically Harvest or Hydro-Rake

- Can not be used on plants which reproduce vegetatively by fragmentation (e.g., milfoil, fanwort, etc.) unless containment can be ensured.
- Can be used only if the waterbody is accessible to machinery.
- Can be used if there is a disposal location available for harvested plant materials.
- Can be used if plant depth is conducive to harvesting capabilities (~ <7 ft. for mower, ~ <12 ft. for hydro-rake).
- Funds are available for repeated harvesting activities in that season.
- A navigation channel is required through dense plant growth.

C. Chemical Treatment

- Can be used if application of chemical is conducted in areas where alternative control techniques are not optimum due to depth, current, use, or type of plant.
- Can be used for treatment of exotic plants where fragmentation is a high concern.
- Can be used where species specific treatment is necessary due to the need to manage other plants (rare or endangered that will not be impacted by chemical treatment).
- Can be used if other methods used as first choices in the past have not been effective.
- A licensed applicator should be contacted to inspect the site and make recommendations about the effectiveness of chemical treatment as compared with

other treatments.

D. Restricted Use Areas (per RSA 487:17, II (d))

- Can be used for exotic species only.
- Can be established in an area that effectively restricts use to a small cove, bay, or other such area where navigation, fishing, and other activities may cause fragmentation to occur.
- Can not be used when there are several “patches” of an infestation of exotic aquatic plants throughout a waterbody.
- Can be used as a temporary means of control.

E. Bottom Barrier

- Can be used for exotic or native species.
- Can be used in small areas, preferably less than 10,000 sq. ft.
- Can be used in an area where the current is not likely to cause the displacement of the barrier.
- Can be used early in the season before the plant reaches the surface of the water.
- Can be used in an area to compress plants to allow for clear passage of boat traffic.
- Can be used in an area to compress plants to allow for a clear swimming area.

F. Drawdown

- Can be used if the target plant(s) are susceptible to drawdown control.
- Can be used in an area where bathymetry of the waterbody would be conducive to an adequate level of drawdown to control plant growth, but where extensive deep habits exist for the maintenance of aquatic life such as fish and amphibians.
- Can be used where plants are growing exclusively in shallow waters where a drawdown would leave this area “in the dry” for a suitable period of time (over winter months) to control plant growth.
- Can be used in winter months to avoid encroachment of terrestrial plants into the aquatic system.
- Can be used if it will not significantly impact adjacent or downstream wetland habitats.
- Can be used if spring recharge is sufficient to refill the lake in the spring.
- Can be used in an area where shallow wells would not be significantly impacted.
- Reference RSA211:11 with regards to drawdown statutes.

G. Dredge

- Can be used in conjunction with a scheduled drawdown.
- Can be used if a drawdown is not scheduled, though a hydraulic pumping dredge should be used.

- Can only be used as a last alternative due to the detrimental impacts to environmental and aesthetic values of the waterbody.

H. Biological Control

- Grass carp cannot be used.
- Exotic controls, such as insects, cannot be introduced to control a nuisance plant.
- Research should be conducted on a potential biological control prior to use to determine the extent of host specificity.


APPENDIX B

SUMMARY OF CONTROL PRACTICES USED IN THE STATE OF NEW HAMPSHIRE FOR EXOTIC AQUATIC PLANTS

Restricted Use Areas:

Restricted Use Areas (RUAs) are a regular control option for lakes with small, contained infestations of exotic plants, limited to small patches or embayments. This is often the case in waterbodies with newly-discovered infestations. RUAs restrict access to all recreational activities in a delineated area to minimize plant fragmentation and thereby reduce the spread of milfoil. As an additional method of protection from fragment migration, RUAs are encircled with a shallow net that is suspended vertically in the water column. The net is approximately 1.5-2.0 feet in height. The top of the net is set to extend four inches above the surface of the water, while the remainder is positioned below the surface of the water (see figure below). This configuration prevents the movement of fragments from infested areas to uninfested areas. Due to the size and nature of net construction, there is no impediment to fish migratory patterns or spawning activities.

Schematic of Restricted Use Area Net



Hand-pulling:

When infestations of exotic aquatic plants begin as single scattered stems or small patches, DES biologists SCUBA dive to hand-pull the plants (and DES can train other certified divers to also perform this management practice). Guidelines for determining feasibility and effective for hand-removal are site specific, but generally sparsely populated patches of up to 5' X 5', single stems, or dense small patch up to 2' X 2' are reasonable.

The whole plant including the roots should be removed in this process, while leaving the beneficial native species intact. This technique works best in softer sediments, with shallow rooted species and for smaller, scattered infestation areas. When hand pulling nuisance species, the entire root system and all fragments of the plants must be collected since small root or stem fragments could result in additional growth of the species. The process must be repeated often to control re-growth of the exotic plants. For a new infestation, hand-pulling activities are typically

conducted several times during the first season, with follow-up inspections for the next 2-5 years or until no re-growth is observed. This control practice has proven successful in many waterbodies.

Mechanical Harvesting

The process of mechanical harvesting is conducted by using machines which cut and collect aquatic plants. These machines can cut the plants up to twelve feet below the water surface. The weeds are cut and then collected by the harvester or other separate conveyer-belt driven device where they are stored in the harvester or barge, and then transferred to an upland site.

The advantages of this type of weed control are that cutting and harvesting immediately opens an area such as boat lanes, and it removes the upper portion of the plants. Due to the size of the equipment, mechanical harvesting is limited to water areas of sufficient size and depth. It is important to remember that mechanical harvesting can leave plant fragments in the water, which if not collected, may spread the plant to new areas. Additionally harvesters may impact fish and insect populations in the area by removing them in harvested material. Cutting plant stems too close to the bottom can result in re-suspension of bottom sediments and nutrients. This management option is only recommended when nearly the entire waterbody is infested, and harvesting is needed to open navigation channels through the infested areas.

Benthic Barriers:

When a small infestation of exotic aquatic plants occurs in clusters of growth (generally areas $>5 \text{ ft}^2$), as opposed to scattered stems, a permeable fiberglass screen can be placed over the area of infested lake sediments. The permeable fabric screening allows for gas release from the sediments while effectively blocking sunlight and compressing the plants into the sediment, inhibiting photosynthesis and eventually killing the plant. Occasionally, in some lakes, gas release from the sediments or boating activity cause the uplifting of screening. Benthic barriers can effectively control small infestations of less than approximately 10,000 square feet.

Benthic barriers have two basic applications. These practices are used to cover pioneering infestations and prevent the spread of the plant. Bottom barriers are installed across small portions of lake bottoms infested with invasive aquatic plants. The disadvantage of benthic barriers is their non-selectivity and limitation of cover to less than 10,000 square feet. Additionally, these physical barriers prevent the growth of all vegetation, which is a necessary component of fish and wildlife habitat.

Bottom barriers are attached to the bottom of a water body by re-bar attached to the edges and across the middle of the material. Bottom barriers are transported to the shoreline adjacent to where installation is to occur. They are then cut to fit the treatment site and rolled onto a length of pipe. Divers carry the roll into the water at the start of the treatment site and secure one edge of the material to the lake bottom. The divers then roll out the remainder of the material and continue to secure it to the bottom sediments. This process is repeated until the plants in the treatment are covered.

Bottom barriers are generally considered for small localized areas rather than lakewide application. Bottom barriers provide 100% control of this weed in areas where they are installed. They also provide long-term control. An ongoing maintenance operation is required to inspect the bottom barrier and clear the mats of sediment buildup.

Benthic barriers are not recommended for application in river systems, as flow can easily uplift the barrier.

Targeted Application of Herbicides:

The use of chemicals, such as herbicides, for the control of noxious and nuisance plant species represents one of the most widely known and effective management options available. Herbicide control of invasive aquatic plants is often the first step in a long-term integrated control program. In the last 15 to 20 years the use and review of herbicides has changed significantly in order to accommodate safety, health, and environmental concerns. Currently no herbicide product can be labeled for aquatic use if it has more than a one in a million chance of causing significant harmful effects to human health, wildlife, or the environment. Because of this, the number of effective and U.S. Environmental Protection Agency (EPA) approved herbicides for aquatic weeds are limited. In most cases the cost and time of testing and registration, rather than environmental issues, limits the number of potentially effective compounds.

All herbicide applications in New Hampshire are performed under permits issued by the New Hampshire Department of Agriculture, Division of Markets and Food, Bureau of Pesticide Control.

Two herbicides have been used in New Hampshire for the control of milfoil. Diquat (trade name Reward), the most often-used herbicide, is a contact herbicide that can generally provide one season of control for milfoil. Because this herbicide does not target the root systems, the plants eventually re-grow from established roots.

The second herbicide, 2, 4-D (trade name Navigate or Aqua Kleen), is a systemic herbicide. It is absorbed into the sediments and taken up through the root system, killing both the roots and the plant biomass above the sediments. Label restrictions for aquatic application currently limit its use in New Hampshire to waterbodies with no water intakes, and with no wells adjacent to the shoreline.

The aquatic herbicide SONAR has been used in New Hampshire to control growths of fanwort. The chemical acts by limiting photosynthesis when chlorophyll-a is affected by the active ingredient of the herbicide.

Extended Drawdown

Water drawdown is used for control of some species of aquatic macrophytes. Drawdown requires some type of mechanism to lower water levels, such as dams or water control structures and use is thus limited. It is most effective when the drawdown depth exceeds the depth or invasion level of the target plant species.

In northern areas, drawdown will result in plant and root freezing during the winter for an added degree of control. Drawdown is typically inexpensive and has intermediate effects (2 or more years). However, drawdown can have other environmental effects and interfere with other functions of the water body (e.g. drinking water, recreation, or aesthetics). Drawdown can result in the rapid spread of highly opportunistic annual weed species, which in most cases is the plant that is targeted for control.

Drawdowns have been used in the past for plant control. In theory, the drying of the plants in the summer, or the freezing of the plants in the winter, will eliminate or limit plant growth. However, milfoil often forms a more succulent terrestrial form during drawdown conditions and the succulent form of the plant can remain viable for long periods of time without submergence, making the practice ineffective. This strategy can be used for control of some native plant species.

Dredging

Dredging is a means of physical removal of aquatic plants from the bottom sediments using a floating or land-based dredge. Dredging can create a variety of depth gradients creating multiple plant environments allowing for greater diversity in lakes plant, fish, and wildlife communities. However due to the cost, potential environmental effects, and the problem of sediment disposal, dredging is rarely used for control of aquatic vegetation alone.

Dredging can take place in to fashion, including drawdown followed by mechanical dredging using an excavator, or using a diver-operated suction dredge while the water level remains up.

Biological Control

There are no approved biological controls for submersed exotic aquatic plant at this time in New Hampshire.

REFERENCES

Department of Environmental Services. 2006: 2006 Section 305(b) and 303(d) Consolidated Assessment and Listing Methodology. November 2005. New Hampshire Department of Environmental Services. NHDES-R-WD-05-29. Available at <http://des.nh.gov/WMB/swqa/calm.html>

Halstead, J.M., J. Michaud, S. Hallas-Burt, and J.P. Gibbs. 2003. "An Hedonic Analysis of Effects of a Nonative Invader (*Myriophyllum heterophyllum*) on New Hampshire (USA) Lakefront Properties." *Environmental Management*. 32 (3): 391 – 398

Luken, J.O. and J.W. Thieret. 1997. *Assessment and Management of Plant Invasions*. Springer-Verlag, New York. 324 pages.